

2020 Forecast: Summer Hypoxic Zone Size Northern Gulf of Mexico

R. Eugene Turner¹ and Nancy N. Rabalais¹

Abstract

A hypoxic water mass with oxygen concentrations $\leq 2 \text{ mg l}^{-1}$ forms in bottom waters of the northern Gulf of Mexico continental shelf each year. Nutrients from the Mississippi River watershed, particularly nitrogen and phosphorus, fertilize the Gulf's surface waters to create excessive amounts of algal biomass, whose decomposition in the bottom layer leads to oxygen depletion. The low oxygen conditions in the Gulf's most productive waters stresses organisms and may even cause their death to threaten living resources, including humans depending on the fish, shrimp and crabs caught there. Various models use the May nitrogen load of the Mississippi River as the main driving force to predict the size of this hypoxic zone in late July. Our prediction is based on one of these models.

The June 2020 forecast of the size of the hypoxic zone in the northern Gulf of Mexico for late July 2020 is that it will cover $20,121 \text{ km}^2$ ($7,769 \text{ mi}^2$) of the bottom of the continental shelf off Louisiana and Texas. The 95% confidence interval is that it will be between $18,054$ and $22,089 \text{ km}^2$ ($6,971$ and $8,529 \text{ mi}^2$). This estimate is based on the assumption that there are no significant tropical storms in the two weeks before the monitoring cruise, or during the cruise. If a storm does occur, then the size of the zone is predicted to be 70% of the predicted size without the storm, equivalent to $14,085 \text{ km}^2$ ($5,438 \text{ mi}^2$).

The predicted hypoxic area is slightly less than the size of the land area of New Hampshire ($23,227 \text{ km}^2$) and 146% of the average of $13,829 \text{ km}^2$ ($n = 34$ including years with storms). If the area of hypoxia becomes as large as predicted, then it will be 4 times the size of the Hypoxia Action Plan goal to reduce the zone to less than $5,000 \text{ km}^2$. No reductions in the nitrate loading from the Mississippi River to the Gulf of Mexico have occurred in the last few decades.

Caveats: 1) This prediction discounts the effect of large storm events that temporarily disrupt the physical and biological system attributes promoting the formation of the low oxygen zone in bottom waters; 2) The potential space on the shelf where hypoxia occurs is limited by the bathymetry; 3) The prediction assumes that there will be no abrupt changes in discharge from now through July; and 4) Unusual weather patterns affecting coastal winds, as experienced in 2009, 2011, 2018, and 2019 may reduce the hypoxic zone size to be lower than predicted.

¹Department of Oceanography and Coastal Sciences
Louisiana State University
Baton Rouge, LA 70803
euturne@lsu.edu
nrabal@lsu.edu

Introduction.....	2
Mississippi River Discharge and Nitrogen Loading.....	4
Hypoxic Zone Size.....	5
Prediction for 2020.....	5
Hypoxia Models and Model Accuracy.....	6
Post-cruise Assessment.....	8
Acknowledgments.....	9
References.....	10
Contacts for Further Information.....	11
Appendices.....	12

Introduction

Hypoxic water masses in bottom waters of the northern Gulf of Mexico occur when the oxygen concentration falls below 2 mg l^{-1} . This hypoxic water is distributed across the Louisiana shelf west of the Mississippi River and onto the upper Texas coast, from near shore to as far as 125 km offshore, and in water depths up to 60 m (Rabalais et al. 2007; Figures 1 and 2). It has been found in all months, but is most persistent and severe in spring and summer (Turner et al. 2005; Rabalais et al. 2007). The July distribution of hypoxic waters most often is a single continuous zone along the Louisiana and adjacent Texas shelf. Hypoxia also occurs east of the Mississippi River delta, but covers less area and is ephemeral. These areas are sometimes called ‘dead zones’ in the popular press because of the absence of commercial quantities of shrimp and fish in the bottom layer – something that is of economic consequence to the fishery (Purcell et al. 2017; Smith et al. 2017). The number of dead zones throughout the world has been increasing in the last several decades and currently totals over 500 (Díaz and Rosenberg 2008; Rabalais et al. 2010; Conley et al. 2011; Breitburg et al. 2018). The dead zone off the Louisiana coast is the second largest human-caused coastal hypoxic area in the global ocean.




Figure 1. The frequency of mid-summer hypoxia (oxygen $\leq 2 \text{ mg l}^{-1}$) over the 70 to 90 station grid on the Louisiana and Texas shelf during the summer from 1985 to 2014. The frequency is determined from those stations for which there are data for at least half of all cruises. From Rabalais et al. (2010).




Figure 2. Oxygen concentrations in bottom water across the Louisiana shelf from July 23 – July 30, 2019. Data source: N.N. Rabalais, Louisiana State University, Louisiana Universities Marine Consortium, and R.E. Turner, LSU; funded by NOAA, National Centers for Coastal Ocean Science.

Systematic mapping of the area of hypoxia in bottom waters of the northern Gulf began in 1985 at geographically fixed stations (Appendix Figure 1). Its size from 1985 to 2019 ranged between 40 to 22,720 km² during late July to early August, and averaged 13,829 km² (5,339 mi²). There were no cruises in 1989 and 2016, and the area was incompletely mapped in 2017. There are few comparable coastwide data for other months, and bi-monthly monitoring on two transects off Terrebonne Bay, LA, and the Atchafalaya delta, LA, ended in 2012. The number of cruises peaked 20 years ago and is now at the bare minimum (Appendix Figure 2).

Hypoxic water masses form from spring to fall on this coast because the consumption of oxygen in bottom water layers exceeds the re-supply of oxygen from the atmosphere. The re-aeration rate is negatively influenced by stratification of the water column, which is primarily dependent on the river's freshwater discharge and accentuated by summer warming. The overwhelming supply of organic matter respired in the bottom layer is from the downward flux of organic matter produced in the surface layer. The transport to the bottom layer is the result of sinking of individual cells, as the excretory products of the grazing predators (zooplankton) that 'package' them as fecal pellets, or as aggregates of cells, detritus and mucus. The respiration of this organic matter declines as it falls through the water column (Turner et al. 1998), but the descent rate is rapid enough that most respiration occurs in the bottom layer and sediments.

The amount of organic matter produced in the surface waters is primarily limited by the supply of nitrogen, not phosphorus (Scavia and Donnelly 2007; Turner and Rabalais 2013), and previous indicators of phosphorus deficiency are not as reliable as they were once thought to be (Fuentes et al. 2014). The evidence for this conclusion is that the supply (loading) of nitrogen (primarily in the form of nitrate-N) from the Mississippi River watershed to the continental shelf within the last few decades is positively related to chlorophyll *a* concentration (Walker and Rabalais 2006; $R^2 = 0.30 - 0.42$), the rate of primary production (Lohrenz et al. 1997, $R^2 > 0.77$;

Lohrenz et al. 2008), and the spatial extent of the hypoxic area in summer (Turner et al. 2012; $R^2 > 0.9$). The size of the shelfwide hypoxic zone has increased since it began occurring in the 1970s simultaneously with 1) the rise in carbon sequestration in sediments, 2) indicators of increased diatom production, and 3) shifts in benthic foraminiferal communities (Turner and Rabalais 1994; Sen Gupta et al. 1996; Turner et al. 2008). There is, therefore, a series of cause-and-effect arguments linking nitrogen loading in the river to phytoplankton production, bottom water oxygen demand, and the formation and maintenance of the largest hypoxic zone in the western Atlantic Ocean.


The oxygen consumption creates a zone of hypoxia that is constrained by the geomorphology of the shelf, horizontal water movement, stratification and vertical mixing (Obenour et al. 2012; Justić and Wang 2014). The significance of reducing nutrient loads to these coastal waters rests on the coupling between the organic matter produced in response to these nutrients and its respiration in the bottom layer (MRNGoM HTF 2001, 2008; Rabalais et al. 2002, 2007, 2010; SAB 2007). The primary driver of the increased nutrient loading is agricultural land use (Alexander et al. 2008; Broussard et al. 2009), which is strongly influenced by farm subsidies (Broussard et al. 2012). The size of the nutrient loading from the river has remained the same in recent decades, or is increasing (Sprague et al. 2011; Crawford et al. 2019).

Mississippi River Discharge and Nitrogen Loading

Hypoxic conditions are dependent on river discharge because of the influence that water volume and salinity have on the physical structure of the water column and on the nutrient load delivered to the coastal zone. The nutrient load is dependent on the: 1) concentration of nutrients, primarily nitrogen, and 2) river discharge. River discharge is, therefore, a key environmental parameter of interest.

The Mississippi River watershed daily discharge in May 2020 was $41,900 \text{ m}^3 \text{ s}^{-1}$ (cms) which is the 48th largest in 52 years from 1968 to 2020, and equal to about 145% of the average May discharge (29,132 cms).

The US Geological Survey (USGS) publishes monthly estimates of nitrogen loading and other aspects of water quality from the Mississippi River watershed into the Gulf of Mexico (<http://toxics.usgs.gov/hypoxia/mississippi/>). The USGS provides information on the data calculations, including an estimate of the 95% confidence range for the nitrogen load. The May nitrite+nitrate (NO_{2+3}) and total nitrogen (TN) load for the Mississippi River watershed for May



is in Figure 3. The total nitrogen load in May is increasingly dominated by the nitrite+nitrate load, which is now about 70% of the TN load.

Figure 3. The annual nitrite+nitrate (NO_{2+3}) and total nitrogen (TN) load for the Mississippi River watershed for May. The estimates are from the USGS.


Hypoxic Zone Size

Models for predicting the size of the hypoxic zone rely on July cruise data primarily because there are no comparable shelfwide data for other months. Data on the size of the hypoxic zone in late July from 1985 to 2019 are based on annual field measurements (data available at <http://www.gulfhypoxia.net>). The 2020 mapping cruise is scheduled for July 25 to August 1. The data will be posted daily at the same web site. There are no values for 1989 (no funding available) or for 2016 (incompatible ship with mechanical breakdown); data from 2017 were incomplete at the end of some transects; data for 1978 to 1984 are estimated from contemporary field data. The estimates for before 1978 assume that there was no significant hypoxia then and are based on results from various models and sediment core analyses. Data for 9 years were not included in the analysis because there were strong storms or unusual wind conditions just before or during the cruise (1998, 2003, 2005, 2008, 2010, 2011, 2013, 2018 and 2019). These storms or unusual wind conditions, by comparison of pre-cruise and post-cruise sampling to data collected during the cruise, changed currents, disrupted the stratified water column, and re-aerated the water column. It may take a few days to several weeks, depending on water temperature and initial dissolved oxygen concentration, for respiration to reduce the dissolved oxygen concentration to $\leq 2 \text{ mg l}^{-1}$ after the water column stratification is re-established. The average reduction in hypoxia size in years with storms compared to years without storms is $70 \pm 9\%$.

Prediction for 2020

We used several models to forecast the hypoxic zone in the northern Gulf of Mexico in July 2020. The most accurate model prediction, we think, is that it will cover $20,121 \text{ km}^2$ ($7,769 \text{ mi}^2$) of the bottom of the continental shelf off Louisiana and Texas. The 95% confidence interval is that it will be between $18,054$ and $22,089 \text{ km}^2$ ($6,971$ and $8,529 \text{ mi}^2$) (Figure 4). This estimate is based on the assumption that there are no significant tropical storms occurring in the two weeks before the monitoring cruise, or during the cruise. If a storm does occur, then the size of the zone is predicted to be 70% of the predicted size without the storm, equivalent to $14,085 \text{ km}^2$ ($5,438 \text{ mi}^2$). This ‘non-storm’ estimate is 46% higher than the average of $13,829 \text{ km}^2$ measured for all years from 1985 to 2019, and the 28th largest measured ($n=34$) since systematic sampling began in 1985.

Figure 4. The measured and estimated size of the hypoxic zone 1979 to 2019 and the predicted for 2020.




The discharge of the Mississippi River has been relatively high this spring, and the Bonnet Carré spillway above New Orleans was opened once, but not the Morganza Spillway. The Bonnet Carré diverts riverwater into Lake Pontchartrain on the east bank, and the Morganza Spillway is used to increase riverflow to the west, through the Atchafalaya Swamp and into the Gulf of Mexico near Morgan City. The movement of water to the east through Bonnet Carré, will probably have a minimal effect on the eastern end of the study area, unlike in 2019 when massive algae blooms occurred in coastal Mississippi and many business shut down.

Hypoxia Models and Model Accuracy

We use several models to predict the size of the hypoxic zone in July that are based on the May total nitrite+nitrate nitrogen load (note: concentration \times discharge equals the nitrite+nitrate load) to the Gulf from the main stem of the Mississippi River and the Atchafalaya River. The residence time of the surface waters along this coast is about 2 to 3 months in the summer, hence the 2 to 3 month lag between the loading rate calculated in May and the size of the hypoxic zone in late July. The stability of these models, however, is not fixed, because the ecosystem is evolving. For example, the size of the hypoxic zone for the same amount of nitrogen loading (as nitrite+nitrate) is increasing at about three times higher over the last 40 years (Figure 5; Turner et al. 2008, 2012). The nitrite+nitrate loading will be referred to here as “nitrate” loading, because the nitrite component is a minimal component of the two. Further, the models will eventually be adjusted to account for the limited space on the shelf for hypoxia to occur (a physiographic constraint). The rapidly developing process-based ecosystem models are a platform to greatly expand understanding how the physical and biological factors interact over all months (Justić and Wang 2014; Justić et al. 2017), are increasingly accurate, are visually appealing, and require additional data to validate them as conditions change throughout the year.

Figure 5. The size of the hypoxic zone per May nitrite+nitrate loading. All years, including strong storm years, are included.




The unstated hypothesis implied by these models is that the system can be treated as a chemostat limited by N, in the same way that the chlorophyll *a* concentration or algal biomass in lakes be modeled by P loading to the lake. The Streeter–Phelps type models initiated by Scavia and colleagues also incorporate this nutrient dose : response framework (Scavia et al. 2003, 2004; Scavia and Donnelly 2007) in their predictive schemes. These models assume that the size of the zone is driven mostly by what happens in the current year and that other influences cause variation around a relatively stable baseline suite of factors. An example of secondary influences might be seasonal or annual variations in wind speed and direction or freshwater volume. Our model is based on the nitrate load of only the current year. The reference point for calibrating the model is the behavior of the system in recent history. We use the last several years of data on the relationship between hypoxic zone size and nutrient loading for this model. Others do something similar. The USGS uses the last five years of data to calibrate the

‘LOADSET’ model, for example, and Scavia and Donnelly (2007) update the coefficients in their model annually by using rolling 3- to 5-year averages for coefficients (Evans and Scavia 2010). Their recent numerical adaptation has the effect of adjusting model input with each year, but not explaining the biological/physical basis for these changes any better than one of our earlier models did with the ‘year’ term. The year term in our model is, in other words, descriptive, but not explanatory beyond the simple nitrogen loading = oxygen deficit relationship.

The estimate for 2020 is in Figure 6. The nitrate data were transformed into their log10 equivalents to avoid the problem encountered in 2012 when the prediction was much larger than the actual size, which is attributable to using a simple linear regression analysis to fit a curvilinear relationship. If there is significant curvature (bowed downward) without this transformation, then both the lower and upper ends of the data field are overestimated. This effect is more dramatic when the relationship is being extended into a sparse data field at the extremes of nitrogen loading, as happened during 2012, which was a drought year with low nitrate loading.

Figure 6. The relationship between nitrate+nitrate in May and the predicted size of the hypoxic zone in July. Several intervals are broken out, with the last one (2000 to 2015) being fit to a regression model. The predicted size of the hypoxic zone for 2020 is indicated by the red dot.



Some of the sensitivity to nitrate loading is carried over from one interval to the next to create a ‘legacy’ effect that may last decades. A legacy effect can be explained as the result of incremental changes in organic matter accumulated in the sediments one year, and metabolized in later years (Turner and Rabalais 1994), by changes in the percent nitrate of the total nitrogen pool (e.g., Figure 6), or by long-term temperature changes (Turner et al. 2017).

Our statistical models, and their predecessors, are fairly accurate models based on past performance (Turner et al. 2008, 2012). The predictions in 2006, 2007, and 2010, for example, were 99%, 107%, and 99%, respectively, of the measured size. The model used here describes 94% of the variation since 2000 (non-storm years). The equivalent model for the Baltic Sea low oxygen conditions explains 49 to 52% of the inter-annual variations in bottom-water oxygen concentration (Conley et al. 2007).

Nutrient load models are robust for long-term management purposes, but they are less robust when short-term weather patterns move water masses or mix up the water column (Rabalais et al. 2018). The size of the hypoxic zone this year is expected to follow the relationship with nitrogen loading—as long as there is no ‘wildcard’ in the form, for example, of

a tropical storm at the time of the annual summer cruise. Some of the variations in the size of the Gulf hypoxic zone result from re-aeration of the water column during storms. The size of the summer hypoxic zone in 2008, for example, was less than predicted because of the influence of Hurricane Dolly. Tropical Storm Don was a similar complication in 2011. Climate changes may alter the spring initiation of hypoxia formation, duration and frequency. The timing of hypoxia in the Chesapeake Bay, for example, is earlier with climate warming (Testa et al. 2018). The needed detailed seasonal data necessary to make phenological comparison are not known. The long-term trend for the northern Gulf of Mexico is that the area of hypoxia is larger for the same amount of nitrogen loading (Turner et al. 2012; Fig. 6).

The prediction in 2018 was noteworthy for the great disparity between the much larger size of the hypoxic zone predicted by *all* models and the actual size. The predicted size of the forecast from four models ranged from 12,949 to 17,523 km², but the measured size was 7040 km². The post-cruise model of the hypoxic zone size revealed a strong change in wind patterns at the time of the cruise. This wind field change resulted in a short-lived decrease in water column stratification and then oxygenation of the bottom layer, particularly on the eastern end of the mapped area.

Other models predicting oxygen dynamics on this shelf are in Bierman et al. (1994), Justić et al. (2003), Scavia and Donnelly (2007), Forest et al. (2011), Scavia et al. (2003, 2004), Testa et al. (2017), Laurent et al. (2018) and Fennel and Laurent (2018). The two other forecasts for this year will be from the University of Michigan (<http://scavia.seas.umich.edu/hypoxia-forecasts/>), Dalhousie University (<http://memg.ocean.dal.ca/news/2020forecast.html>), and the Virginia Institute of Marine Science (http://www.vims.edu/research/topics/dead_zones/forecasts/gom/index.php). The NOAA ensemble predictions are based on these models (<http://www.noaa.gov/media-releases>). These models do not always produce similar results, and model improvement is one focus of ongoing research efforts supported by the NOAA National Centers for Coastal Ocean Science. The general result from an ensemble analysis using the four model results indicates that a 60% reduction in Mississippi River nitrogen load is required to reach the Hypoxia Task Force goal, and that a 25% load reduction is required to have a 95% certainty of observing a hypoxic area reduction within a consecutive 5-year assessment period (Scavia et al. 2017).

The data from this year's cruise will be used to quantify the relative merits of the assumptions of the models, and to compare them with other models. This is an example of how long-term observations are one of the best ways to test and calibrate ecosystem models, to recognize the dynamic nature of our changing environment(s), and to improve the basis for sound management decisions.

A general recent description of hypoxia in the northern Gulf of Mexico is in Rabalais and Turner (2019). We review the past, present, and possible future conditions of hypoxia in the northern Gulf of Mexico and provide some insight into possible management actions.

Post-cruise Assessment

A post-cruise assessment will be provided at the end of the summer shelfwide hypoxia cruise and posted on the same website where this report appears (<http://www.gulfhypoxia.net>).

Acknowledgments

Support is from the NOAA National Centers for Coastal Ocean Science. The hypoxia area data are from <http://www.gulfhypoxia.net>. We thank John Wickham of NOAA and Brent T. Aulenbach, Mike Woodside and colleagues at the USGS for providing the nitrogen loading data for the Mississippi River.

References

- Alexander, R.B., R.A. Smith, G.E. Schwarz 2008. Differences in phosphorus and nitrogen delivery to the Gulf of Mexico from the Mississippi River basin. **Environmental Science and Technology** 42: 822–830.
- Bierman, V.J., Jr., S.C. Hinz, D. Zhu, W.J. Wiseman, Jr., N.N. Rabalais, R.E. Turner 1994. A preliminary mass balance model of primary productivity and dissolved oxygen in the Mississippi River plume/inner Gulf shelf region. **Estuaries** 17: 886–899.
- Breitburg, D., L.A. Levin, A. Oschlies, M. Grégoire, F.P. Chavez, D.J. Conley, V. Garçon, D. Gilbert, D. Gutiérrez, K. Isensee, G.S. Jacinto, K.E. Limburg, I. Montes, S.W.A. Naqvi, G.C. Pitcher, N.N. Rabalais, M.R. Roman, K.A. Rose, B.A. Seibel, M. Telszewski, M. Yasuhara, J. Zhang 2018. Declining oxygen in the global ocean and coastal waters. **Science** 359: eaam7240, 11 pp.
- Broussard, W., R.E. Turner 2009. A century of changing land use and water quality relationships in the continental U.S. **Frontiers in Ecology and the Environment** 7: 302-307.
- Broussard, W., R.E. Turner, J. Westra 2012. Do federal farm policies and agricultural landscapes influence surface water quality? **Agriculture, Ecosystems & Environment** 158: 103-109. 10.1016/j.agee.2012.05.022
- Conley, D.J., J. Carstensen, G. Ærtebjerg, P.B. Christensen, T. Dalsgaard, J.L.S. Hansen, A. B. Josefson 2007. Long-term changes and impacts of hypoxia in Danish coastal waters. **Ecological Applications** Supp. 17: S165-S184.
- Conley, D.J., 18 co-authors 2011. Hypoxia is increasing in the coastal zone of the Baltic Sea. **Environmental Science and Technology** 45: 6777–6783. doi.org/10.1021/es201212r.
- Crawford, J.T., E.G. Stets, L.A. Stets 2019. Network controls on mean and variance of nitrate loads from the Mississippi River to the Gulf of Mexico. **Journal of Environmental Quality** 48:1789-1799.
- Díaz, R.J., R. Rosenberg 2008. Spreading dead zones and consequences for marine ecosystems. **Science** 321: 926-929.
- Evans, M.A., D. Scavia 2010. Forecasting hypoxia in the Chesapeake Bay and Gulf of Mexico: Model accuracy, precision, and sensitivity to ecosystem change. **Environmental Research Letters** 6: 015001.
- Fennel, K., A. Laurent 2018. N and P as ultimate and proximate limiting nutrients in the northern Gulf of Mexico: implications for hypoxia reduction strategies, **Biogeosciences** 15: 3121-3131.
- Justić, D., L. Wang 2014. Assessing temporal and spatial variability of hypoxia over the inner Louisiana-upper Texas shelf: Application of an unstructured-grid three-dimensional coupled hydrodynamic-water quality model. **Continental Shelf Research** 72: 163-179.
- Justić, D., N.N. Rabalais, R.E. Turner 2003. Simulated responses of the Gulf of Mexico hypoxia to variations in climate and anthropogenic nutrient loading. **Journal of Marine Systems** 42: 115-126.
- Justić, D., K.A. Rose, R.D. Hetland, K. Fennel (Eds.) 2017. Modeling Coastal Hypoxia: Numerical Simulations of Patterns, Controls and Effects of Dissolved Oxygen Dynamics.


- Springer, NY. 433 pp.
- Laurent, A., K. Fennel, D.S. Ko, J. Lehrter 2018. Climate change projected to exacerbate impacts of coastal eutrophication in the northern Gulf of Mexico. **Journal of Geophysical Research-Oceans** 123: 3408–3426. doi: 10.1002/2017JC013583 (2018)
- Lohrenz, S.E., G.L. Fahnenstiel, D.G. Redalje, G.A. Lang, X. Chen, M.J. Dagg 1997. Variations in primary production of northern Gulf of Mexico continental shelf waters linked to nutrient inputs from the Mississippi River. **Marine Ecology Progress Series** 155: 45–54.
- Lohrenz, S.E., D.G. Redalje, W.J. Cai, J. Acker, M.J. Dagg 2008. A retrospective analysis of nutrients and phytoplankton productivity in the Mississippi River Plume. **Continental Shelf Research** 28: 1466–1475.
- Maiti, V., S. Fang, J. Guinness, N.N. Rabalais, J. Craig, D.R. Obenour 2018. A space-time geostatistical assessment of hypoxia in the northern Gulf of Mexico, **Environmental Science & Technology** 52: 12484-12493 doi: 10.1021/acs.est.8b03474
- MRNGoM HTF (Mississippi River Nutrient/Gulf of Mexico Hypoxia Task Force) 2001. *Action Plan for Reducing, Mitigating, and Controlling Hypoxia in the Northern Gulf of Mexico*; Office of Wetlands, Oceans, and Watersheds, U.S. Environmental Protection Agency; Washington, DC.
- MRNGoM HTF (Mississippi River Nutrient/Gulf of Mexico Hypoxia Task Force) 2008. *Action Plan for Reducing, Mitigating, and Controlling Hypoxia in the Northern Gulf of Mexico*. Office of Wetlands, Oceans, and Watersheds, U.S. Environmental Protection Agency, Washington, D.C.
- Obenour, D.R., A.M. Michalak, Y. Zhou, D. Scavia 2012. Quantifying the impacts of stratification and nutrient loading on hypoxia in the northern Gulf of Mexico. **Environmental Science and Technology** 46: 5489–5496.
- Purcell, K.M., J.K. Craig, J.M. Nance, M.D. Smith, L.S. Benneer 2017. Fleet behavior is responsive to a large-scale environmental disturbance: Hypoxia effects on the spatial dynamics of the northern Gulf of Mexico shrimp fishery. **PLoS ONE** 12(8): e0183032. <https://doi.org/10.1371/journal.pone.0183032>
- Rabalais, N.N., R.E. Turner, D. Scavia 2002. Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi River. **BioScience** 52: 129-142.
- Rabalais, N.N., R.E. Turner, B.K. Sen Gupta, D.F. Boesch, P. Chapman, M.C. Murrell 2007. Hypoxia in the northern Gulf of Mexico: Does the science support the plan to reduce, mitigate and control hypoxia? **Estuaries and Coasts** 30: 753-772.
- Rabalais, N.N., R.J. Díaz, L.A. Levin, R.E. Turner, D. Gilbert, J. Zhang 2010. Dynamics and distribution of natural and human-caused hypoxia. **Biogeosciences** 7: 585-619.
- Rabalais, N.N., L.M. Smith, R E. Turner. 2018. The *Deepwater Horizon* oil spill and Gulf of Mexico shelf hypoxia. **Continental Shelf Research** 152: 98-107.
- Rabalais, N.N., R.E. Turner 2019. Gulf of Mexico Hypoxia: Past, present and future. **Limnology and Oceanography Bulletin** 28: 117-124. doi.org/10.1002/lob.10351
- Scavia, D., N.N. Rabalais, R.E. Turner 2003. Predicting the response of Gulf of Mexico hypoxia to variations in Mississippi River nitrogen load. **Limnology and Oceanography** 48: 951-956.
- Scavia, D., D. Justić, V.J. Bierman, Jr. 2004. Reducing hypoxia in the Gulf of Mexico: Advice from three models. **Estuaries** 27: 419–425.
- Scavia, D., K.A. Donnelly 2007. Reassessing hypoxia forecasts for the Gulf of Mexico. **Environmental Science and Technology** 41: 8111-8117.
- Scavia, D., I. Bertani, D.R. Obenour, R.E. Turner, D.R. Forrest, A. Katkin 2017. Ensemble

- modeling informs hypoxia management in the northern Gulf of Mexico. **Proceeding National Academy of Sciences (USA)** 114: 8823–8828. doi/10.1073/pnas.1705293114
- Science Advisory Board (SAB) 2007. Hypoxia in the northern Gulf of Mexico, An Update. U.S. Environmental Protection Agency, Science Advisory Board (SAB) Hypoxia Panel Advisory, Report EPA-SAB-08-003, Environmental Protection Agency, Washington, D.C.
[http://yosemite.epa.gov/sab/sabproduct.nsf/C3D2F27094E03F90852573B800601D93/\\$File/EPA-SAB-08-003complete.unsigned.pdf](http://yosemite.epa.gov/sab/sabproduct.nsf/C3D2F27094E03F90852573B800601D93/$File/EPA-SAB-08-003complete.unsigned.pdf)
- Sen Gupta, B.K., R.E. Turner, N.N. Rabalais 1996. Seasonal oxygen depletion in continental-shelf waters of Louisiana: Historical record of benthic foraminifers. **Geology** 24: 227-230.
- Smith, M.D., A. Oglen, A.J. Kirkpatrick, F. Asche, L.S. Benneer, J.K. Craig, J.M. Nance 2017. Seafood prices reveal impacts of a major ecological disturbance. **Proceedings National Academy of Sciences (USA)** 114: 1512–1517, doi: 10.1073/pnas.1617948114
- Sprague, L.A., R.M. Hirsch, B.T. Aulenbach 2011. Nitrate in the Mississippi River and its tributaries, 1980 to 2008: Are we making progress? **Environmental Science and Technology** 45: 7209-7216. dx.doi.org/10.1021/es201221s
- Testa, J.M., J.B. Clark, W.C. Dennison, E.C. Donovan, A.W. Fisher, W. Ni, M. Parker, D. Scavia S.E. Spitzer, A.M. Waldrop, V.M.D. Vargas, G. Ziegler 2017. Ecological forecasting and the science of hypoxia in Chesapeake Bay. **BioScience** 67: 614–626.
- Testa, J.M., R.R., Murphy, D.C. Brady, W.M. Kemp, 2018. Nutrient- and climate-induced shifts in the phenology of linked biogeochemical cycles in a temperate estuary. **Frontiers in Marine Science** 5: 114. doi: 10.3389/fmars.2018.00114
- Turner, R.E., N. Qureshi, N.N. Rabalais, Q. Dortch, D. Justić, R. Shaw, J. Cope 1998. Fluctuating silicate:nitrate ratios and coastal plankton food webs. **Proceedings National Academy of Sciences (USA)** 95: 13048-13051.
- Turner, R.E., N.N. Rabalais 1994. Coastal eutrophication near the Mississippi river delta. **Nature** 368: 619-621
- Turner, R.E., N.N. Rabalais, E.M. Swenson, M. Kasprzak, T. Romaire 2005. Summer hypoxia, Northern Gulf of Mexico: 1978 to 1995. **Marine Environmental Research** 59: 6577.
- Turner, R.E., N.N. Rabalais, D. Justić 2008. Gulf of Mexico hypoxia: Alternate states and a legacy. **Environmental Science and Technology** 42: 2323-2327.
- Turner, R.E., N.N. Rabalais, D. Justić 2012. Predicting summer hypoxia in the northern Gulf of Mexico: Redux. **Marine Pollution Bulletin** 64: 318-323. DOI: 10.1016/j.marpolbul.2011.11.008
- Turner, R.E., N.N. Rabalais, D. Justić 2017. Trends in summer bottom-water temperatures on the Northern Gulf of Mexico continental shelf from 1985 to 2015. **PloS One** 12(9): e0184350. <https://doi.org/10.1371/journal.pone.0184350>
- Walker, N.D., N.N. Rabalais 2006. Relationships among satellite chlorophyll a, river inputs, and hypoxia on the Louisiana continental shelf, Gulf of Mexico. **Estuaries and Coasts** 29: 1081–1093.


Contacts for Further Information

Nancy N. Rabalais (LSU, 225-578-8531 wk, 985-870-4203 c, 985-851-2836 wk);
nrabalais@lumcon.edu)
 R. Eugene Turner (LSU, eturne@lsu.edu)


Appendix



Appendix Figure 1. Location of hypoxia monitoring stations sampled in summer (not every year, depending on location of hypoxic area), the transects off Terrebonne Bay (transect C) and Atchafalaya Bay (transect F), and the ocean observing system (asterisk) off Terrebonne Bay (no longer in operation).



Appendix Figure 2. The number of State, Federal and university cruises associated with hypoxia measurements in the northern Gulf of Mexico from 1985 to 2016. LUMCON = Louisiana Universities Marine Consortium; SEAMAP = Southeast Area Monitoring and Assessment Program; TAMU = Texas A&M University; UMCES = University of Maryland Center for Environmental Studies; EPA = U.S. Environmental Protection Agency; NECOP = Nutrient Enhanced Coastal Ocean Productivity; LDWR = Louisiana Department of Wildlife Research; LUMCON-T = transects sampled during the year by LUMCON. Source: Maiti et al. 2018; used with permission.



Appendix Figure 3. The daily river discharge at Tarbert Landing, LA, from 1935 through 31 May 2020. Units are cubic feet per second \times 1000. Figure modified from <http://rivergages.mvr.usace.army.mil/WaterControl/Districts/MVN/tar.gif>.